

# 2016 SCIENTIFIC REPORT



MINISTRY OF FOOD AND DRUG SAFETY

National Institute  
of Food and Drug Safety Evaluation

---

## Risk Assessment of Omethoate

Omethoate, a metabolite of dimethoate, is an organophosphorous pesticide. Its MRL in Korea is set at 0.01–3 mg/kg for 39 foodstuffs including potato and celery (MRLs for Pesticides in Foods, May 31, 2016).

The ADI of omethoate at 0.0003 mg/kg bw/day was established by applying the safety factor 100 (differences between species and individual entities) to the NOAEL of 0.03 mg/kg bw/day drawn from the multi-generation study on rats.

The intake amount of omethoate was estimated based on the results of the analysis of 3,516 samples of 65 foodstuffs including rice in the Monitoring of Agricultural Products in Korea (2006–2009) by the National Institute of Food and Drug Safety Evaluation. The result of the monitoring showed that the pesticide level was below the LOQ, and thus, omethoate was not detected in any of the samples. Concerning data lower than the LOQ, in case more than 60% of data were below the LOQ, estimation was made by applying 0 (non-detection) as the lower exposure limit or LOQ (upper exposure limit), according to the “evaluation of low level contamination of foods” recommended by the WHO. Food consumption was calculated through SAS 9.4 using the tertiary food code data from the KNHANES conducted for five years (2010–2014). For the average weight of all age groups, 60 kg, the weight currently (as of 2016) being applied for establishment and revision of pesticide residue standards, was used. Risk characterization was made by calculating the HI in consideration of the EDI calculated in the exposure assessment and the ADI, the safe level of human exposure.

In general, when HI is 1 or higher, the adverse effects of toxicity are expected from the exposure, and when HI is lower than 1, adverse effect is not expected. The results of the risk assessment of omethoate in all age groups revealed HI between 0 (non-detection data 0

# 2016 SCIENTIFIC REPORT



MINISTRY OF FOOD AND DRUG SAFETY

National Institute  
of Food and Drug Safety Evaluation

applied) and 0.043 (non-detection data LOQ applied), as shown in the table below, and that its concentration is within the safe level of human exposure.

**Table 1.** ADI and HI of omethoate

Age	EDI (mg/person/day)		Average weight (kg)	EDI (mg/kg bw/day)		ADI (mg/kg bw/day)	HI	
	0	LOQ (mg/kg)		0	LOQ (mg/kg)		0	LOQ (mg/kg)
All	0	$0.8 \times 10^{-3}$	60	0	$0.1 \times 10^{-4}$	0.0003	0	0.043

**Key words:** Omethoate, Risk Assessment, Organophosphorus insecticide, ADI, Monitoring, Pesticide